Determination of the Postmortem Interval - between Biological Indicators and Forensic Interpretation

Alina Daniela PĂTRU*

Abstract

The determination of the postmortem interval represents a fundamental component of forensic investigation, particularly in cases where establishing the timeline of events is essential for reconstructing the circumstances of death. This study examines the relationship between biological indicators of postmortem change and their interpretation within a forensic context, emphasizing the need for a correlated and applied approach. Biological processes such as body cooling, lividity, and rigor mortis provide measurable data that may assist in estimating the time since death. In practice, these indicators are influenced by multiple variables, including environmental conditions, body position, clothing, and contact with surrounding surfaces. For example, a body exposed to lower temperatures may exhibit delayed postmortem changes, while a body in a warmer environment may show accelerated progression. Such variations affect the reliability of estimation when indicators are considered in isolation. The analysis highlights the importance of correlating biological findings with scene-related elements, including the position of the body, trace evidence, and environmental characteristics. The postmortem interval is not established through a single parameter, but through the integration of multiple sources of information that must be interpreted together. By approaching the postmortem interval as both a biological and interpretative construct, the study underlines the need for caution and methodological rigor. Estimation must remain grounded in observable data while accounting for the variability inherent in postmortem processes.

Keywords: *postmortem interval; algor mortis; livor mortis; rigor mortis; forensic interpretation*

1. Introduction

Establishing the postmortem interval is a central task in forensic investigation, as it provides the temporal framework necessary for reconstructing the sequence of

* Assistant Professor, PhD, Spiru Haret University, Romania, Corresponding author: alina.patru@spiruharet.ro.

Copyright: © 2026 by the authors.
Open access publication under the terms and conditions of the
Creative Commons Attribution (CC BY) license
https://creativecommons.org/licenses/by/4.0/

events. Determining when death occurred allows investigators to correlate physical evidence, evaluate statements, and place the victim within a specific timeframe. In many cases, this estimation becomes critical when other sources of information are limited or absent. The process relies primarily on the observation of postmortem changes that occur in the body after death. Body cooling, lividity, and rigor mortis represent the most frequently used indicators. These processes follow general patterns, yet their progression is influenced by the conditions in which the body is found. Temperature, humidity, clothing, and contact with surrounding surfaces all affect the rate and expression of postmortem changes.

In practice, these indicators rarely provide precise values when considered individually. For example, a body found in a cold environment may retain heat longer than expected, while one exposed to higher temperatures may cool more rapidly. Similarly, lividity patterns may indicate the position of the body after death, but they may also reveal whether the body has been moved. Such observations demonstrate that biological indicators must be interpreted in relation to the context in which they appear. The determination of the postmortem interval therefore extends beyond the simple measurement of biological processes. It requires the correlation of multiple elements, including scene characteristics and other forensic findings. The position of the body, the presence of traces, and environmental conditions all contribute to the estimation process. Within this framework, the postmortem interval is not a fixed or exact value, but an approximation based on observable data and contextual analysis. Its accuracy depends on the ability to integrate biological indicators with the conditions of the scene, while accounting for the variability that characterizes postmortem processes.

2. Biological Indicators of the Postmortem Interval

The estimation of the postmortem interval is primarily based on the analysis of biological changes that occur in the body after death. Among these, body cooling, lividity, and rigor mortis are the most commonly used indicators. Each of them provides useful information, but only when interpreted in relation to the conditions in which the body is found.

Body cooling, or algor mortis, begins immediately after death as the body gradually loses heat and approaches the ambient temperature. In controlled conditions, this process follows a relatively predictable pattern. In practice, however, the rate of cooling varies significantly. A body located in a cold environment, exposed to wind or moisture, may cool faster than expected, while a body in a warm or enclosed space may retain heat for a longer period. Clothing, body mass, and contact with surfaces also influence heat loss. For example, a body lying on a cold surface may exhibit uneven cooling, with certain areas losing heat more rapidly than others.

Lividity, or livor mortis, results from the settling of blood in the lower parts of the body under the influence of gravity. It becomes visible as discoloration of the skin

and may provide information about body position after death. In practice, its distribution is particularly relevant. If lividity is observed on surfaces that do not correspond to the position in which the body is found, this may indicate that the body has been moved. For instance, discoloration on the back combined with a body found in a face-down position suggests a change in position after death. The degree of fixation of lividity also contributes to estimation, as it indicates how long the blood has remained in a particular position (Drăgan, 2014, p. 118).

Rigor mortis, the stiffening of muscles after death, develops progressively and affects different muscle groups in sequence. Its presence and distribution may provide additional information regarding the elapsed time since death. In practice, however, its progression is influenced by factors such as temperature, physical condition, and the circumstances of death. A body exposed to higher temperatures may develop rigor more rapidly, while a body in a cooler environment may show delayed onset. Similarly, intense physical activity prior to death may accelerate its development. These indicators do not function independently. Their value emerges when they are considered together. For example, a body that is still warm, with no visible lividity and no signs of rigor, suggests a short interval since death. In contrast, a cold body with fixed lividity and fully developed rigor indicates a longer interval. Intermediate situations require careful correlation, as the indicators may not align perfectly due to environmental or situational factors.

In applied forensic analysis, the interpretation of these biological changes must remain flexible. Each case presents a specific combination of conditions that influence how postmortem processes develop. For this reason, estimation cannot rely on a single indicator, but must be based on the integration of all observable elements within the context of the scene.

3. Environmental and External Factors Influencing the Postmortem Interval

The progression of postmortem changes is not determined solely by biological processes, but is significantly influenced by the environment in which the body is found. In practical forensic work, these external factors often explain discrepancies between expected and observed postmortem indicators. Temperature represents the most influential factor. Low temperatures slow down biological processes, delaying the onset and progression of rigor mortis and reducing the rate of body cooling. Conversely, high temperatures accelerate decomposition and may lead to a rapid development and disappearance of rigor. For example, a body found outdoors during winter may retain characteristics associated with an earlier postmortem stage, while a body exposed to heat may appear more advanced than the actual interval would suggest.

Humidity and moisture also play an important role. A body found in a dry environment may undergo dehydration, leading to preservation effects such as

mummification, which slow down decomposition. In contrast, a body exposed to moisture or submerged in water may exhibit accelerated tissue breakdown. Water environments introduce additional variables, including temperature fluctuations and biological activity, which further complicate estimation. Contact with surfaces influences both cooling and lividity. A body lying on a cold or conductive surface may cool unevenly, while pressure points may prevent the formation of lividity in certain areas. These aspects become relevant when interpreting the distribution of postmortem changes, particularly in relation to body position (Moise & Stancu, 2023, p. 78).

Clothing and coverings also affect the rate of change. Layers of clothing may retain heat, delaying cooling and altering the progression of rigor mortis. At the same time, clothing may influence the visibility and distribution of lividity, as well as the preservation of certain areas of the body. The location of the body, whether indoors or outdoors, introduces further differences. Indoor environments tend to provide more stable conditions, while outdoor settings expose the body to variable temperatures, weather conditions, and biological activity. Insects and other environmental agents may accelerate decomposition, especially in warmer conditions, contributing additional indicators but also increasing variability.

These factors demonstrate that postmortem indicators cannot be interpreted independently of their context. A given biological sign may have different significance depending on the environment in which it is observed. For this reason, estimation of the postmortem interval requires not only the observation of biological changes, but also the careful evaluation of external conditions. In applied forensic practice, discrepancies between indicators are often explained by environmental influences rather than by errors in observation. Understanding how these factors interact with biological processes is essential for producing a more accurate and context-based estimation.

4. Correlation of Postmortem Indicators in Forensic Interpretation

The estimation of the postmortem interval becomes meaningful only when biological indicators and environmental factors are examined together. In practice, no single parameter provides a reliable estimation on its own. The interpretative process depends on the correlation of multiple elements, each contributing partial information that must be integrated into a coherent assessment.

In many cases, indicators do not align perfectly. A body may exhibit partial rigor mortis while also presenting signs of advanced cooling, or lividity may appear inconsistent with the position in which the body is found. Such situations require analysis rather than reliance on standard expectations. For example, a cold body with incomplete rigor may initially suggest a longer postmortem interval; however, if the body was exposed to low temperatures, the delayed development of rigor becomes a relevant factor in interpretation (Grecu, 2025a, p. 132).

The correlation of lividity with body position represents another important aspect. If lividity is fixed in areas that do not correspond to the current position of the body, this may indicate postmortem movement. When this observation is combined with trace evidence, such as disturbed surfaces or signs of handling, the interpretation becomes more consistent and allows for a clearer reconstruction of events. Body cooling must also be evaluated in context. A measured body temperature may suggest a certain interval since death, but without considering environmental conditions, this estimation may be misleading. For instance, a body found in a warm indoor environment may retain heat longer, while one exposed to cold air currents may cool more rapidly than expected (Grecu, 2025a, p. 133).

The process of correlation often involves resolving apparent contradictions. Instead of treating discrepancies as errors, they must be examined as potential indicators of specific conditions or actions. In this sense, inconsistency between indicators may itself provide useful information, pointing to environmental influences or postmortem changes in body position. In applied forensic analysis, estimation is therefore constructed through comparison and adjustment. Each indicator is evaluated in relation to the others, and conclusions are drawn based on the degree of consistency between them. Where multiple elements support the same timeframe, the estimation gains reliability. Where such consistency is lacking, the interpretation remains approximate and must be expressed accordingly.

The postmortem interval should thus be understood as the result of a correlated analysis rather than a direct measurement. Its determination depends on the ability to integrate biological data, environmental conditions, and scene-related observations into a structured and context-based interpretation.

5. Practical Implications in Forensic Investigation

In forensic investigation, the estimation of the postmortem interval plays a central role in structuring the analysis of events and correlating available evidence. Its practical value becomes evident when it is used not as an isolated indicator, but as a reference point for evaluating other elements within the case.

One of the most relevant applications concerns the verification of timelines. The estimated time since death may be compared with witness statements, reported activities, or declared alibis. When discrepancies arise, they do not automatically invalidate one source of information, but require careful reassessment. For example, a shorter postmortem interval than expected may suggest that the timeline of events needs to be reconsidered, while a longer interval may indicate either an earlier time of death or the influence of environmental conditions on postmortem changes. The position and condition of the body also have direct practical implications. Lividity patterns that do not correspond to the current position of the body may indicate postmortem movement. When such findings are correlated with other elements, such as disturbed surfaces, drag marks, or the absence of expected traces, the

interpretation gains consistency. In this way, the postmortem interval is not analyzed separately, but contributes to reconstructing actions that occurred after death (Pătru, 2024, p. 98).

Another important application involves the assessment of scene integrity. The stage of postmortem changes may indicate whether the body has remained undisturbed for a certain period or whether external intervention has occurred. For instance, inconsistencies between the expected development of rigor mortis and the observed condition of the body may suggest that environmental conditions have changed or that the body has been repositioned. The estimation of the postmortem interval also influences investigative priorities. In cases where death is recent, the scene may still preserve transient traces that require immediate documentation (Grecu, 2025b, p. 183). In contrast, when a longer interval is indicated, the focus may shift toward identifying persistent evidence and reconstructing earlier stages of the event. The condition of the body, therefore, guides not only interpretation, but also the strategy of evidence collection.

In more complex situations, such as advanced decomposition, the role of the postmortem interval becomes more interpretative than descriptive. Biological indicators may be less precise, requiring greater reliance on environmental data, scene context, and additional forensic findings. In such cases, estimation is expressed in broader intervals rather than exact values.

In applied forensic work, the usefulness of the postmortem interval lies in its capacity to support correlation rather than to provide definitive answers. It functions as a temporal framework that allows investigators to organize evidence, identify inconsistencies, and evaluate possible sequences of events. Its relevance depends on how effectively it is integrated with other findings within the overall analytical process.

6. Conclusion

Estimating the postmortem interval does not result from a single observation, but from the combination of multiple indicators that must be interpreted together. Body cooling, lividity, and rigor mortis provide useful reference points, yet their significance becomes clear only when they are examined in relation to the conditions in which the body is found. In practice, environmental factors frequently explain differences between expected and observed postmortem changes. Temperature, humidity, and body position may alter the rate and appearance of these processes, making direct estimation unreliable if context is ignored. For this reason, biological indicators must always be evaluated alongside scene-related elements (Grecu, 2025a, p. 94).

Situations in which indicators do not align are common. A body may present partial rigor while also showing advanced cooling, or lividity may not correspond to the

current position. Such inconsistencies should not be treated as errors, but as information that may reflect movement of the body or the influence of external conditions. Interpreting these aspects is part of the analytical process. The postmortem interval is therefore best understood as an approximation that supports the reconstruction of events rather than as an exact measurement. Its reliability increases when multiple elements converge and decreases when evidence is limited or altered. In forensic investigation, its value lies in how it is used. When integrated with other findings, it helps define the temporal framework of the case, supports the evaluation of statements, and contributes to identifying inconsistencies. A consistent interpretation depends on correlation, not on isolated indicators.

Approaching the postmortem interval in this manner allows the analysis to remain grounded in observable data while addressing the variability of postmortem processes. The focus shifts from precise calculation to structured interpretation, where conclusions reflect both the available evidence and the conditions under which it developed.

7. References

Drăgan, J. (2014). *Criminalistică: Manual de instrucție judiciară/ Criminalistics: A Manual of Judicial Investigation*. Bucharest: Romania de Mâine Foundation.

Grecu, P. (2025). Impact of Environmental Conditions on the Persistence and Quality of Fingermarks: Implications for Crime Scene Investigation. *Journal of Law and Public Administration*, Volume XI, Issue 22, pp. 132–139.

Grecu, P. (2025a). Investigating Death by Asphyxia – Correlating Scene Findings with Autopsy Results. *Current Issues in Law and Socio-Economic Relations. Scientific Journal*, Issue 1(7), Volume 1, Kiev, Ukraine, pp. 92–99.

Grecu, P. (2025b). The Significance of Defensive Injuries in Interpreting the Mechanism of Assault. *International Scientific and Practical Conference "Expert Support for Organized Crime Investigation: International Experience and National Practice"*, Odessa, Ukraine, November 27, 2025, pp. 182–185.

Moise, A. C., & Stancu, E. (2023). *Elemente metodologice de investigare a infracțiunilor: Curs universitar/ Methodological Elements for the Investigation of Crimes: University Course* (2nd ed.). Bucharest: Universul Juridic.

Pătru, A. D. (2024). *Criminalistică/Criminalistics*. Craiova: Sitech.